

ISSN-0873-0122 (Print) + ISSN-0873-0124 (Electronic)

Volume 15 / Number 2 / April-June 2012



Indian Journal of Forensic Medicine & Toxicology

Website: www.ijfml.com

Official Organ of Indian Association of Medical Legal Experts (IAMLE)

Link Jurnal : <https://medicopublication.com/index.php/ijfml/article/view/17984>

Link Pdf artikel : <https://medicopublication.com/index.php/ijfml/article/view/17984/15803>

Home / Editorial Team

Editorial Team

Editor in Chief

Prof S K Dhattarwal

Forensic Medicine, PGIMS, Rohtak, Haryana
E-mail: editor_ijfmt@gmail.com

Editor

Dr. T. Nataraja Moorthy

Forensic Science Faculty of Health & Life Sciences ,Management & Science University, Malaysia

EDITORIAL ADVISORY BOARD

1. Prof Sudhir K Gupta, Head, FMT, AIIMS, New Delhi, India
2. Prof Mete Gulmen ,Cukurova University, TURKEY
3. Prof. Leandro Duarte De Carvalho , Minas Gerais, Belo Horizonte, BRAZIL
4. Dr. Valery Gunas, National Pirogov Memorial Medical University, Vinnytsya, UKRAINE
5. Dr. Rahul Pathak Forensic Science, Dept of Life Sciences ,Anglia Ruskin University, Cambridge, UNITED KINGDOM
6. Prof Emilio Nuzzalese, University of Turin, Italy
7. Dr Noha A. Magdy Elrafie, Forensic Toxicology, Ain Shams University, Cairo, EGYPT
8. Dr Rituja Sharma, Associate Prof, Law, Banasthali Vidyapeeth Jaipur
9. Dr Shankar Bakkanwar (Associate Professor) Forensic Medicine, Kasturba Medical College, Manipal, Karnataka
10. Dr K. Ravikumar , Raksha Shakti University, Ahmedabad, Gujarat.
11. Dr C. Behera (Asst. Prof) Dept of FMT, AIIMS, New Delhi
12. Dr. Kanak Lata Verma, Deputy Director, Toxicology, RPSL, Chanakyapuri New Delhi
13. Dr. Asha Srivastava (Senior Scientific Officer) Forensic Psychology, Central Forensic Science Laboratory, CBI, Delhi
14. Dr. Raghendra Kumar Vidua, (Associate Prof), FMT, AIIMS Bhopal
15. Dr. Valbhav Saran (Asst.Prof.) School of Forensic Science, Sam Higginbottom Institute of Agriculture Technology & Sciences, Allahabad
16. Ms Aparna R. Asst. Prof. Forensic Serology & Biology, Jain University, Bengaluru
17. Dr. Deepali Jain, Asst Prof, Forensic Science, JBB Ambedkar University, Lucknow
18. Prof. NK Aggrawal Forensic Medicine, UCMs, Delhi
19. Prof. Manoj Kumar Mohanty, Forensic Medicine, AIIMS, Bhubneshwar, Odisha
20. Prof. Amar Jyoti Patowary, Forensic Medicine, NEIGRIHMS, Shillong, Meghalaya
21. Prof S. Venkata Raghava - Forensic Medicine, Bangalore Medical College, Bengaluru
22. Prof. Shalini Gupta Oral Pathology and Microbiology, Faculty of Dental Sciences, King George Medical University, Lucknow
23. Prof. Virender Kumar Chhoker Forensic Medicine, Santosh Medical College, Ghaziabad, UP
24. Prof. Dayanand G Gannur , Forensic Medicine , Shri BM Patil Medical College, Hospital & Research centre, Bijapur, Karnataka
25. Prof Praveen Arora, Forensic Medicine, SAIMS, Indore, MP
26. Prof Barkha Gupta , Saraswathi Institute of Medical Sciences Hapur, Uttar Pradesh India
27. Prof M Prashant Apollo Medical College Hyderabad
28. Prof Dimple Patel , Anatomy, AMC MET Medical College , Ahmedabad , Gujarat
29. Dr Mohammed Nader Shalaby, Associate Professor of Biological Sciences and Sports Health Department, Faculty of Physical Education, Suez Canal University, Egypt

Current Issue



[Make a Submission](#)

Browse

[Open Journal Systems](#)

Information

[For Readers](#)

[For Authors](#)

[For Librarians](#)

Home / Archives / Vol. 16 No. 2 (2022): Indian Journal of Forensic Medicine & Toxicology

Vol. 16 No. 2 (2022): Indian Journal of Forensic Medicine & Toxicology



DOI: <https://doi.org/10.37506/ijfmt.v16i2>

Published: 2022-04-12

Current Issue



[Make a Submission](#)

Browse

[Open Journal Systems](#)

Information

[For Readers](#)
[For Authors](#)
[For Librarians](#)

Articles

Comparative Study between Variable Computer Printed Documents to Identify the Type of the Producing Output Devices

Abeer Sayed Abdo Borae, Yasser Fouad Abd El-Monem Al-Akid, Khaled Abd El-Azeem Mahmoud Tamim, Nesrine Abd El-Rahman Mahmoud 1-10



A Correlation Study between Fingerprints and Lip Prints among Twins

M.Sugatha, R.Jenifer Jacqueline, M.Taqiuddinkhan 11-19



The Relation between the Plasma Level of Testosterone Hormone and the Severity of COVID-19 in Iraqi Patients

Rusul A. Abass, Manal K Rasheed, Ghassan T. Saeed 326-333



The Effects of Water Provision and Education on Students' Hydration Status, Cognitive Abilities, and Fine Motor Function in A Full-Day Primary School

Saidah Rauf, Santi Aprilian Lestaluhi 334-342



Effect of Lifestyle Modification Guidelines on Mild Preeclampsia

Sally Ebrahim Rashad Ali, Ragaa Ali Mohamed Abdraboo, Nagat Salah Shalaby, Inaam Hassan Abdelati 343-350



Dhikr and Prayer Guidance on Peace of Mind and Blood Pressure Control

Samsualam, Masriadi 351-357



Emotional and Cognitive Impairment among Cardiac Patients - A Narrative Review

Sarika M L, Sasmita Das, Suresh Kumar Behera 358-365



The Effects of Water Provision and Education on Students' Hydration Status, Cognitive Abilities, and Fine Motor Function in A Full-Day Primary School

Saidah Rauf¹, Santi Aprilian Lestaluhu²

¹Masohi Nursing Study Program, Nursing Departement, Politeknik Kesehatan Kemenkes Maluku, Masohi, Indonesia, ²Nutrition, Departement, Politeknik Kesehatan Kemenkes Maluku, Ambon, Indonesia

Abstracts

Background/aim: Dehydration due to inadequate water intake in children may disrupt brain functions. This study aimed to investigate the effects of hydration education with water provision on hydration status, cognitive functions and motor performance.

Materials and Methods: Forty-eight children in primary full-day school (age 8 – 11 years) were recruited in this study. The intervention of hydration education was performed for two days, both for students and teachers. The water was provision for 12 days. Letter cancellation task (LCT), symbol digit modalities test (SDMT), and direct image different test (DIDT) were used to assess cognitive functions. Motor performance was evaluated using a finger tapping test (FTT). Hydration status was determined with urine specific gravity (USG) and urine color (UC). Before and after interventions, all parameters were evaluated.

Results: Most students found dehydrated (>65%). The students had lower USG and UC after the intervention and it was correlated with each other (before; $r_s=0.45$, $P = 0.001$ and after; $r_s=0.82$, $P = 0.001$). All cognitive and motor functions were significantly higher after an intervention ($P < 0.05$).

Conclusion: Hydration education to students and teachers combined with water provision at primary full-day school increases hydration status and enhances cognitive and fine motor skills.

Keywords: Urine color, Attention, Motor function, Dehydration

Background

Growing evidence shows an increasingly widespread lack of total fluid intake, mostly water,

among children worldwide⁽¹⁻⁴⁾. Children and adolescents (aged 4–17) drank less than 500 mL of water daily^(3,4). Besides, the fact that water is the most abundant part of the body of children places them at a high risk of dehydration^(5,6).

Corresponding author:

Saidah Rauf

Masohi Nursing Study Program, Nursing
Departement, PoltekkesKemenkes Maluku, Masohi,
Indonesia Masohi, 97511, Indonesia

Email addresses: saidahrauf@poltekkes-maluku.ac.id

Increased awareness of the inadequate water intake induces dehydration of school-aged children, and access to water in schools has been raised⁽⁷⁾. Stookey and Konig found significant variations in fluid availability for school-aged children in Mexico,

Brazil, Argentina, China, and Indonesia. In some states, more than half of students do not have free access to drinking water while in school⁽⁸⁾, particularly in hot and dry areas⁽⁷⁾. According to Kausik et al., promoting adequate school water intake can help students' biological functioning⁽⁹⁾.

Several investigations discovered that dehydration could impair cognitive function^(10,11). In addition, education intervention⁽¹²⁾ and water supplementation were sufficient to prevent dehydration^(11,13). Acute water supplementation increases the student's visual focus and motor abilities⁽¹³⁾. Therefore, the effect of combining intervention (education and water provision) on cognitive and fine motor functions have not been explicitly studied in primary full-day school.

The current study examined the effects of hydration education and water provision for 12 days on urine hydration markers (USG and UC), cognitive functions (LCT, SDMT, and DIDT), and fine motor performance (FTT). We used all parameters as a baseline before the intervention. The degree of change in each parameter was measured after the intervention.

Methods

Participants

This study was carried out in September 2019 in Masohi, Central Maluku, Indonesia. Before collecting data, primary school principals were asked for permission. Selection criteria included grade (4-6) and health. Chronic disease or medication use were excluded. Their teachers informed their parents about the research. Before data collection, participants and their guardians signed consent forms. Only 48 of the 58 participants with parental consent completed the study. In this study, the Helsinki Declaration's ethical principles were followed (1964). All PoltekkesKemenkes Maluku procedures were approved by the Ethical Committee (LB.02.02/6.2/2910/2019).

Study design and education intervention

The study was quasi-experimental. All participants received hydration education and water provision. The research had three stages. Pre-intervention assessments included hydration, cognition, and fine motor. This lasted only one day. After that, teachers and participants learned about daily fluid requirements, dehydration symptoms, and rehydration options. Participants' water intake was monitored for two weeks by trained teachers who also drank water at school. Participants were observed at school from 7:00 am to 4:00 pm. Each class got refillable water. A 500 mL water bottle from home was also required. All school activities resume as scheduled. On the last day, we did post-intervention. Pre-intervention parameters were assessed.

Hydration status assessment

Pre- and post-intervention hydration status was assessed between 3:30 and 4:00 p.m. using USG and UC. Each participant received one sterile urine container. USG was measured twice with a portable refractometer (Cole-Parmer RSA-BR82T). Between samples, the refractometer was calibrated with distilled water. The sample was placed in a transparent container against a white background to calculate UC using a validated eight-color scale.

Cognitive and motor assessments

Letter cancellation task

The LTC requires participants to locate and cross a target letter within a grid of distractor letters. This test measured visual attention. This study used the one-letter version. Subjects had to cancel 'U' between O and V letters. Participants had 60 seconds to hit all targets. Scores were calculated by subtracting errors from the number of correct targets.

Symbol digit modalities test

The SDMT's cognitive demand is on visual information processing speed and attention. The top of the paper sheet has nine stimulus symbols and nine Arabic numbers from 1 to 9. Above the key were 120 blank boxes with a single pattern. Initial practice (5 min) for each participant, with orders. Participants have 90 seconds to fill in the blank boxes with the key. The number of empty squares that could be filled with the correct symbol determined the score.

Direct image difference test

This test was for visual attention. Two similar images were shown side by side. Participants had 60 seconds to spot the differences between the two images. In this case, more correct differences found meant better results (maximum =12).

Finger tapping test

Fine motor assessment was done by tapping digitally (CNS-Finger Tapping Test, Tushar Kalra). Participants had 60 seconds to practice with the instrument before the test. The digital app records several taps in a 3-trial by a preferred hand (10 s for each). On the digital screen until the trial is finished. The fine motor score was the average of three taps.

Statistical Analysis

The statistical analyses were performed using GraphPad Prism 9.0.0 and SPSS version 21 software programs. The data were presented in means \pm standard deviation (SD) or median \pm interquartile range (IQR) based on the normality distribution. Significance levels were set at $p < 0.05$.

The participants' characteristics were analyzed between males and females using unpaired t-test or Mann-Whitney test, and Kolmogorov-Smirnov test. The mean difference of participants' scores on FTT was analyzed with paired t-test. The mean rank differences of UC, USG, and cognitive assessments test pre and post-intervention using Wilcoxon test.

The differences of cognitive and motor assessments between hydration status were analyzed using one-way ANOVA or Kruskal Wallis test. Spearman correlation test was used to assess the correlation between US and USG levels.

Results

Participant characteristics

Table 1 shows participant characteristics for boys and girls. The Kolmogorov-Smirnov test revealed that both boys and girls had similar school grades ($p > 0.05$). Age, weight, BMI, and hydration status were not different between genders ($p > 0.05$). Only 22% of boys and 33% of girls had well-hydrated urine color samples. According to USG, one-third of boys and one-quarter of girls were severely dehydrated (≥ 1.030).

Hydration status based on urinary markers

Figure 1 presents data on the effect of hydration education and water provision at class on participants' urine hydration markers. There was a significant difference in the UC ($p < 0.0001$) and USG ($p < 0.0001$) between pre (Fig. 2a) and post interventions (Fig. 2b). Both urine hydration marker values were lower after intervention than that before the intervention.

Spearman correlation test showed a significantly positive correlation between the US and USG levels on pre and post-intervention ($p = 0.001$). The correlation coefficient between the US and USG levels before treatment with hydration education and water provision was moderate ($r_s = 0.45$), while after treatment was intense ($r_s = 0.82$).

Cognitive and motor performances

The results of LCT, SDMT, DIDT, and FTT before and after treatment are presented in Figure 2. There were significant differences in the mean rank of LCT, SDMT, and DIDT between pre and post-

intervention ($p < 0.0001$). All cognitive performances improved after treatment. The paired t-test also found significant differences in the mean of FTT between the time of assessments ($p < 0.0001$).

The cognitive and motor performances pre and post-intervention were also analyzed based on hydration status, specifically USG value (Table 2). The LCT values pre and post-treatment with hydration education and water provision were significantly differences (one-way ANOVA, $F_{3, 44} = 3.84$; $p = 0.016$ and $F_{3, 44} = 2.86$; $p = 0.048$, respectively). Post-hoc analyses of these data revealed that the significant dehydration group showed a significant decrease in LCT than well-hydrated ($p = 0.003$ and $p = 0.005$, pre and post) and minimal dehydration ($p = 0.008$

and $p = 0.040$, pre, and post) groups. Interestingly, there was no significant decrease in LCT scores on the serious dehydration group than the well-hydrated group, both pre and post-intervention.

The SDMT mean rank did not change significantly between pre- and post-treatment ($p > 0.005$). Dehydration also reduces DIDT score post-intervention ($p = 0.041$). The FTT data showed a significant difference between groups post-intervention (one-way ANOVA, $F_{3, 44} = 6.99$; $p = 0.001$) but not pre-intervention. The serious dehydration group had significantly fewer FTT scores than the well-hydrated ($p < 0.0001$), minimal dehydrated ($p < 0.0001$), and significant dehydrated ($p = 0.001$) groups.

Table 1 Characteristics of participants between male and female

Characteristics	Boy (n = 27)	Girl (n=21)	P value
Age (years) ^a	10 ± 3	9 ± 1	0.135*
8 – 9	12 (44%)	12 (57%)	
10 – 11	15 (56%)	9 (43%)	
Grades at school			
4th grade	11 (41%)	13 (62%)	0.665#
5th grade	9 (33%)	7 (33%)	
6th grade	8 (26%)	1 (5%)	
Body weight (Kg) ^b	28.3 ± 14	27.5 ± 10.5	0.442*
BMI-b	-0.28 ± 1.7	0.54 ± 0.9	0.435 [†]
Underweight	3 (11%)	0	
Normalweight	16 (59%)	21 (100%)	
Overweight	4 (15%)	0	
Obese	4 (15%)	0	

^aData presented as mean ± SD. ^bData presented as median ± IQR.

*Mann-Whitney test. #Kolmogorov-Smirnov test. [†]Unpaired t-test. n=number of participant.

Table 2. Cognitive and fine motor performances of participants in pre and post intervention based on USG

Cognitive and motor assessments		Hydration status based on USG				p value
		WH	MD	SiD	SeD	
	n Pre/post	15/36	13/16	6/4	14/2	
LCT	Pre	29.8 ± 4.4	28.9 ± 6.2	20.5 ± 9.3*#	25.7 ± 6.4	0.016a
	Post	33.1 ± 4.6	32.3 ± 3.8	25.6 ± 7.9*#	33.0 ± 2.8	0.046a
SDMT	Pre	21 ± 25	13 ± 24	17 ± 12.3	20 ± 11.3	0.426b
	Post	25 ± 9.8	30.5 ± 19.8	25.3 ± 6.3	19	0.207b
DIDT	Pre	7 ± 3	5 ± 2	6 ± 2	5 ± 3	0.041b
	Post	7 ± 3	8.5 ± 1	8.5 ± 6	5	0.161b
FTT	Pre	51.3 ± 6.9	48.8 ± 8.9	48.3 ± 6.2	48.6 ± 4.9	0.685a
	Post	53.4 ± 5.5	52.2 ± 3.1	52 ± 4.5	41.5 ± 3.5*#§	0.026a

^aThe data were analyzed using One-way Anova. ^bThe data were analyzed using Kruskalwallis test. *<0.05 vs. WH ;#<0.05 vs. MD ; §<0.05 vs. SiD. WH, Well-hydrated; MD, Mild dehydration; SiD, Significant dehydration; SeD, Serious dehydration; LCT, Letter cancelation task; SDMT, Symbol digit modalities test (SDMT); DIDT, Direct image difference test; FTT, Finger tapping test.




Figure 1

The effect of hydration education and water provision of drinking water at class on participants’ urine hydration markers. a. Urine color. b. Urine specific gravity. The data are presented as median±IQR (n=51).

****p<0.0001 (Wilcoxon test).rs, Spearman rank correlation coefficient. p, p-value




Figure 2

The effect of hydration education on cognitive and motor performances of participants. **a. Cognitive assesments.** The data are presented as median±IQR (n=48). ****p<0.0001 (Wilcoxon test). **b. Fine motor assesment.** The data are presented as mean±SD (n=48). ****p<0.0001 (Paired t-test).

LCT, Letter cancelation task; SDMT, Symbol digit modalities test (SDMT); DIDT, Direct image difference test; FTT, Finger tapping test.

The effect of hydration education and water provision of drinking water at class on participants' urine hydration markers. **a. Urine color.** **b. Urine specific gravity.** The data are presented as median±IQR (n=51). ****p<0.0001 (Wilcoxon test). r_s , Spearman rank correlation coefficient. *p*, *p*-value

Discussion

The current study found that two weeks of hydration education and water provision improved students' hydration and cognitive abilities in primary full-day school. Improved hydration status (lower USG and UC values) influenced cognitive and fine motor performance based on dehydration levels.

Our study had shown that the incidence of dehydration before the intervention was high (68% - 73%). This research finding was consistent with previous studies^(10,14,15). These findings show that once children have reached school, they do not drink enough water⁽¹³⁾. As a result, the risk of dehydration increases with extended schooling time.

The hydration status of boys and girls did not differ statistically (Table 1). Some experiments had varying results. They found that boys' urine osmolality and USG were higher than girls⁽¹⁶⁻¹⁹⁾. The reason for our result remains uncertain at present since we did not perform a drinking diary or food recall, but it may be linked to variations in the race or ethnicity of participants⁽²⁰⁾ in this study compared to those previous reports⁽¹⁶⁻¹⁹⁾.

The current study found that after two weeks of intervention, USG and UC decreased significantly (p<0.05), indicating an increase in school-based water intake. It has been proposed that increasing water consumption is more important than other drinks in reducing dehydration⁽²⁰⁾. It is well-known that providing water at school increases student water intake and hydration^(11,13,15,21). Water access and verbal hydration education reduced urine osmolality, USG, and UC in children attending sports camps⁽¹²⁾. However, teachers' role as role models and monitoring

student hydration seems to have helped improve students' hydration status. According to Schätzer et al., healthy school hydration initiatives should help empower teachers⁽²²⁾. A lower urine osmolality or adequate hydration were found in schools that provided water, drinking education, and structured drinking agreements⁽²³⁾. This study's approach to reducing child dehydration includes both hydration education and teacher empowerment.

It has been conveyed that a loss of 2% of body weight due to inadequate hydration could detract cognitive and psychomotor performances⁽²⁴⁾. Kozioł-Kozakowska and colleagues reported that dehydration was demonstrated as a risk factor (Odds Ratio= 2.85) for impairment of students' attention⁽²⁵⁾. These dehydration effects on brain function have been restored during water intake⁽²⁶⁾. Along with improved hydration, the study found improved cognitive function on the LCT, SDMT, and DIDT scores (Fig. 2a) and fine motor abilities FTT (Fig. 2b). The LCT, SDMT, and DIDT are recognized for determining visual attention⁽¹⁴⁾. Our results matched those of previous studies^(15,21,27,28).

As shown in Table 2, there were statistically significant ($p < 0.05$) differences in mean LCT scores between groups of hydration status. Also, the SDMT scores were identical in both studies. Pre-intervention DIDT test scores showed a significant difference between hydration status groups, whereas post-intervention FTT scores showed no difference. This result suggests that hydration status specifically affected cognitive performances. Previous research found that body water adequacy was domain-specific rather than generalized^(21,29).

The group with the worst attention scores on LCT pre- and post-intervention was significantly dehydrated ($p < 0.05$), according to post hoc analysis (Table 2). Surprisingly, the LCT score was similar in both groups ($p > 0.05$). The severe dehydration

group also had shallow fine motor performances on FTT post-intervention. Extreme dehydration affected children's fine motor skills ($p > 0.05$), indicating that dehydration affects children's fine motor skills. Previously, studies evaluating dehydration-related mental performance showed a dose-response relationship. The more body fluids are lost because of dehydration, the worse the performance will be^(30,31). Our FTT results support these viewpoints, but not for LCT scores. In children with extreme dehydration, the attention is well-preserved on LCT after the intervention is possibly attributable to physiological mechanisms of adaptation. They were experienced extreme and severe dehydration since the pre-intervention period. Physiological systems respond acutely to hypohydration to reduce homeostatic disruption and probably adjust to chronic water deficit to increase tolerance⁽³²⁾.

Conclusion

Hydration education to student and teacher combined with water provision for 12 days at primary full-day school may return student USG and UC to well-hydrated levels. Improving the hydration status of students has been shown to enhance cognitive and motor skills. Our study's limitation does not consider the calculation of the daily amount of intake of water or other beverages and provided a control group. Further investigations on the effects of hydration education with water supplementation on different aspects or parameters of children's brain function are also warranted.

Source of Funding This study was funded by the PoltekkesKemenkes Maluku under contract number YM/03.02/1/5219/2019. This grant was awarded to authors S. Rauf and S. A. Lestaluhu.

Conflict of Interest Nil

References

1. Gouda Z, Zarea M, El-Hennawy U, Viltard M,

- Lepicard E, Hawili N, et al. Hydration Deficit in 9- to 11-Year-Old Egyptian Children. *Glob Pediatr Heal*. 2015;2:2333794X1561178.
2. Iglesia I, Guelinckx I, De Miguel-Etayo PM, González-Gil EM, Salas-Salvadó J, Kavouras SA, et al. Total fluid intake of children and adolescents: cross-sectional surveys in 13 countries worldwide. *Eur J Nutr*. 2015;54(2):57-67.
 3. Guelinckx I, Iglesia I, Bottin JH, De Miguel-Etayo P, González-Gil EM, Salas-Salvadó J, et al. Intake of water and beverages of children and adolescents in 13 countries. *Eur J Nutr*. 2015;54:69-79.
 4. Gandy J, Martinez H, Carmuega E, Arredondo JL, Pimentel C, Moreno LA, et al. Fluid intake of Latin American children and adolescents: results of four 2016 LIQ.IN 7 National Cross-Sectional Surveys. *Eur J Nutr*. 2018;57(3):53-63.
 5. Bottin JH, Morin C, Guelinckx I, Perrier ET. Hydration in Children: What Do We Know and Why Does it Matter? *Ann Nutr Metab*. 2019;74(Suppl3):11-8.
 6. Perrier ET, Armstrong LE, Bottin JH, Clark WF, Dolci A, Guelinckx I, et al. Hydration for health hypothesis: a narrative review of supporting evidence. *Eur J Nutr*. 2020;(0123456789).
 7. Bar-David Y, Ukrin J, Landau D, Bar-Davids, Pilpel D. Voluntary dehydration among elementary school children residing in a hot arid environment. *J Hum Nutr Diet*. 2009;22(5):455-60.
 8. Stookey JD, König J. Describing water intake in six countries: results of Liq.In7 surveys, 2015-2018. *Eur J Nutr*. 2018;57(3):35-42.
 9. Kaushik A, Mullee MA, Bryant TN, Hill CM. A study of the association between children's access to drinking water in primary schools and their fluid intake: Can water be "cool" in school? *Child Care Health Dev*. 2007;33(4):409-15.
 10. Bar-David Y, Ukin J, Kozminsky E. The effect of voluntary dehydration on cognitive functions of elementary school children. *Acta Paediatr Int J Paediatr*. 2005;94(11):1667-73.
 11. Fadda R, Rapinett G, Grathwohl D, Parisi M, Fanari R, Calò CM, et al. Effects of drinking supplementary water at school on cognitive performance in children. *Appetite*. 2012;59(3):730-7.
 12. Kavouras SA, Arnaoutis G, Makrillos M, Garagouni C, Nikolaou E, Chira O, et al. Educational intervention on water intake improves hydration status and enhances exercise performance in athletic youth. *Scand J Med Sci Sport*. 2012;22(5):684-9.
 13. Booth P, Taylor B, Edmonds C. Water supplementation improves visual attention and fine motor skills in school children. *Educ Heal*. 2012;30(3):75-9.
 14. Chard AN, Trinies V, Edmonds CJ, Sogore A, Freeman MC. The impact of water consumption on hydration and cognition among school children: Methods and results from a crossover trial in rural Mali. *PLoS One*. 2019;14(1):1-14.
 15. Trinies V, Chard AN, Mateo T, Freeman MC. Effects of water provision and hydration on cognitive function among primary-school pupils in Zambia: A randomized trial. *PLoS One*. 2016;11(3):1-14.
 16. Kavouras SA, Bougatsas D, Johnson EC, Arnaoutis G, Tsipouridi S, Panagiotakos DB. Water intake and urinary hydration biomarkers in children. *Eur J Clin Nutr*. 2017;71(4):530-5.
 17. Bougatsas D, Arnaoutis G, Panagiotakos DB, Seal AD, Johnson EC, Bottin JH, et al. Fluid consumption pattern and hydration among 8-14 years-old children. *Eur J Clin Nutr*. 2018;72(3):420-7.
 18. Ebner A, Manz F. Sex difference of urinary osmolality in German children. *Am J Nephrol*. 2002;22(4):352-5.
 19. Skinner R, Cole M, Pearson ADJ, Coulthard MG,

- Craft AW. Specificity of pH and osmolality of early morning urine sample in assessing distal renal tubular function in children: Results in healthy children. *Br Med J.* 1996;312(7042):1337–8.
20. Kenney EL, Long MW, Craddock AL, Gortmaker SL. Prevalence of inadequate hydration among US children and disparities by gender and race/ethnicity: National Health and Nutrition Examination Survey, 2009–2012. *Am J Public Health.* 2015;105(8):e113–8.
 21. Khan NA, Westfall DR, Jones AR, Sinn MA, Bottin JH, Perrier ET, et al. A 4-d Water Intake Intervention Increases Hydration and Cognitive Flexibility among. *J Nutr.* 2019;149(12):2255–64.
 22. Schatzer M, Schätzer J, Hoppichler F. School Programs in Austria: A Combined Behavioral and Environmental Intervention to Promote Healthy Hydration. *Ann Nutr Metab.* 2019;74(Suppl3):25–9.
 23. Michels N, Van den Bussche K, Vande Walle J, De Henauw S. School Policy on Drinking and Toilets: Weaknesses and Relation With Children’s Hydration Status. *J Nutr Educ Behav.* 2019;51(1):32–40.
 24. Grandjean AC, Grandjean NR. Dehydration and Cognitive Performance. *J Am Coll Nutr.* 2007;26(5):549S–554S.
 25. Koziół-Kozakowska A, Piórecka B, Suder A, Jagielski P. Body composition and a school day hydration state among polish children—a cross-sectional study. *Int J Environ Res Public Health.* 2020;17(19):1–12.
 26. Biller A, Reuter M, Patenaude B, Homola GA, Breuer F, Bendszus M, et al. Responses of the human brain to mild dehydration and rehydration explored in vivo by 1H-MR imaging and spectroscopy. *Am J Neuroradiol.* 2015;36(12):2277–84.
 27. Perry CS, Rapinett G, Glaser NS, Ghetti S. Hydration status moderates the effects of drinking water on children’s cognitive performance. *Appetite.* 2015;95:520–7.
 28. Tung SEH, Ch’ng YZ, Karnan T V., Chong PN, Zubaidah JO, Chin YS. Fluid intake, hydration status and its association with cognitive function among adolescents in petaling perdana, selangor, malaysia. *Nutr Res Pract.* 2020;14(5):490–500.
 29. Patsalos OC, Thoma V. Water supplementation after dehydration improves judgment and decision-making performance. *Psychol Res.* 2020;84:1223–34.
 30. Gopinathan PM, Pichan G, Sharma VM. Role of dehydration in heat stress-induced variations in mental performance. *Arch Environ Health.* 1988;43(1):15–7.
 31. Adan A. Cognitive performance and dehydration. *J Am Coll Nutr.* 2012;31(2):71–8.
 32. Akerman AP, Tipton M, Minson CT, Cotter JD. Heat stress and dehydration in adapting for performance: Good, bad, both, or neither? *Temperature.* 2016;3(3):412–36.

**LEMBAR
HASIL PENILAIAN SEJAWAT SEBIDANG ATAU PEER REVIEW
KARYA ILMIAH : JURNAL ILMIAH**

Judul Jurnal Ilmiah (Artikel) : The Effects of Water Provision and Education on Students' Hydration Status, Cognitive Abilities, and Fine Motor Function in A Full-Day Primary School

Jumlah penulis : 2 Orang (Saidah Rauf, Santi Aprilian Lestaluhi)

Status Pengusul : Penulis Pendamping

Identitas Jurnal Ilmiah : a. Nama Jurnal : Indian Journal of Forensic Medicine & Toxicologi
 b. Nomor ISSN : p-0973-9122 and e-0973-9130
 c. No/Vol/ Bln/Th : No 2/Vol 16/April 2022
 d. Penerbit : Indian Association of Medico-Legal expert
 e. Jumlah halaman : 9 halaman (334-342)
 f. DOI artikel : <https://doi.org/10.37506/ijfimt.v16i2.17984>

Web jurnal : <https://medicopublication.com/index.php/ijfimt/article/view/17984>

Kategori Publikasi Jurnal Ilmiah : Jurnal Ilmiah Internasional
 (beri ✓ pada kategori yang tepat) Jurnal Ilmiah Nasional
 Jurnal Ilmiah Nasional Terakreditasi
 Jurnal Ilmiah Nasional Tidak Terakreditasi


Hasil Penilaian *Peer Review* :

Komponen Yang Dinilai	Nilai Maksimal Jurnal Ilmiah : 30				Nilai Akhir Yang Diperoleh (40%)
	Internasional <input checked="" type="checkbox"/>	Nasional <input type="checkbox"/>	Nasional Terakreditasi <input type="checkbox"/>	Nasional Tidak Terakreditasi <input type="checkbox"/>	
a. Kelengkapan unsur isi artikel (10%)	3				3
b. Ruang lingkup dan kedalaman pembahasan (30%)	9				8
c. Kecukupan dan kemutahiran data/informasi dan metodologi (30%)	9				7
d. Kelengkapan unsur dan kualitas penerbit (30%)	9				9
Total = (100%)	30				26

Komentar :

Kelengkapan dan unsur isi artikel : **Sangat baik**, Ruang Lingkup dan kedalaman pembahasan : **Baik**, Kecukupan dan kemutahiran data/informasi dan metodologi : **Cukup Baik**. Kelengkapan unsur dan kualitas penerbit : **Sangat Baik**,

Ambon, 18 April 2022
Reviewer 1 (Satu)


 Abd. Rivai S. Dunggio, A.KP., M.Kes
 NIP. 197502051998031001
 Pembina / IV-a/ Lektor Kepala
 Poltekkes Kemenkes Maluku

**LEMBAR
HASIL PENILAIAN SEJAWAT SEBIDANG ATAU PEER REVIEW
KARYA ILMIAH : JURNAL ILMIAH**

Judul Jurnal Ilmiah (Artikel) : The Effects of Water Provision and Education on Students' Hydration Status, Cognitive Abilities, and Fine Motor Function in A Full-Day Primary School

Jumlah penulis : 2 Orang (Saidah Rauf, Santi Aprilian Lestaluhu)

Status Pengusul : Penulis Pendamping

Identitas Jurnal Ilmiah : a. Nama Jurnal : Indian Journal of Forensic Medicine & Toxicologi
b. Nomor ISSN : p-0973-9122 and e-0973-9130
c. No/Vol/ Bln/Th : No 2/Vol 16/April/2022
d. Penerbit : Indian Association of Medico-Legal expert
e. Jumlah halaman : 9 halaman (334-342)
f. DOI artikel : <https://doi.org/10.37506/ijfmt.v16i2.17984>

Web jurnal : <https://medicopublication.com/index.php/ijfmt/article/view/17984>

Kategori Publikasi Jurnal Ilmiah : Jurnal Ilmiah Internasional
(beri ✓ pada kategori yang tepat) Jurnal Ilmiah Nasional
 Jurnal Ilmiah Nasional Terakreditasi
 Jurnal Ilmiah Nasional Tidak Terakreditasi

Hasil Penilaian Peer Review :

Komponen Yang Dinilai	Nilai Maksimal Jurnal Ilmiah : 30				Nilai Akhir Yang Diperoleh (40%)
	Internasional <input checked="" type="checkbox"/>	Nasional <input type="checkbox"/>	Nasional Terakreditasi <input type="checkbox"/>	Nasional Tidak Terakreditasi <input type="checkbox"/>	
a. Kelengkapan unsur isi artikel (10%)	4				3
b. Ruang lingkup dan kedalaman pembahasan (30%)	9				8
c. Kecukupan dan kemutahiran data/informasi dan metodologi (30%)	9				7
d. Kelengkapan unsur dan kualitas penerbit (30%)	9				9
Total = (100%)	30				26

Komentar :

Kelengkapan dan unsur isi artikel : **Sangat baik**, Ruang Lingkup dan kedalaman pembahasan : **Baik**, Kecukupan dan kemutahiran data/informasi dan metodologi : **Cukup Baik**. Kelengkapan unsur dan kualitas penerbit : **Sangat Baik**,

Ambon, 18 April 2022
Reviewer 2 (dua)



Mahmud, S.Pd., M.Kes
NIP. 196108171986031004
Penata Tk.1 / III-d / Lektor
Poltekkes Kemenkes Maluku